

Kähler-Einstein metrics on orbifolds and Einstein metrics on spheres

Alessandro Ghigi and János Kollár

February 2, 2008

Abstract

A construction of Kähler-Einstein metrics using Galois coverings, studied by Arezzo–Ghigi–Pirola, is generalized to orbifolds. By applying it to certain orbifold covers of $\mathbb{CP}^n$ which are trivial set theoretically, one obtains new Einstein metrics on odd-dimensional spheres. The method also gives Kähler-Einstein metrics on degree 2 Del Pezzo surfaces with A_1 or A_2 –singularities.

1 Introduction

The aim of this paper is to explain how the methods of Arezzo, Ghigi, and Pirola [1] can be applied to construct Kähler-Einstein metrics on compact complex orbifolds with positive first Chern class, and then use the approach of Boyer, Galicki, and Kollár [10] to obtain new Einstein metrics on odd dimensional spheres.

The somewhat unusual aspect is that we work with orbifolds $\mathcal{X}$ that admit a map $\pi : \mathcal{X} \rightarrow \mathbb{P}^n$ which is the *identity* map set theoretically. Nonetheless, in the orbifold category π is a nontrivial Galois cover, although with trivial Galois group.

The existence of Kähler-Einstein metrics on compact complex manifolds with positive first Chern class is still a difficult problem. For surfaces and toric manifolds a complete solution is known, due respectively to Tian [25] and Wang-Zhu [28]. Apart from these cases, there are two large classes of examples. The simplest are homogeneous spaces, for instance $\mathbb{P}^n$, quadrics, Grassmannians. In all these cases, the first Chern class is large, meaning for instance, that it is a large multiple of a generator of $H_2(X, \mathbb{Z})$. The opposite case, when the first Chern class is a small multiple of a generator of $H_2(X, \mathbb{Z})$ is also understood in many instances; see [8] for a good overview.

A blending of these two approaches was developed in Arezzo, Ghigi, and Pirola [1] to yield Kähler-Einstein metrics on certain manifolds X which can be realized as Galois covers of another manifold Y with a Kähler-Einstein metric. Since the method relies on finite group actions, it is most successful when symmetries form a natural part of the complex structure, for instance for double covers of $\mathbb{P}^n$.

A construction of Einstein metrics on odd dimensional spheres was studied in Boyer, Galicki, and Kollár [10]. The idea is that the quotient of an odd dimensional sphere by a circle action is frequently a complex orbifold, and a result of Kobayashi [16] allows one to lift a Kähler-Einstein orbifold metric from the quotient to an Einstein metric on the sphere.

A frequently occurring case, studied by Orlik and Wagreich [21] and Boyer, Galicki, and Kollár [10], appears when the quotient S^{2n+1}/S^1 is $\mathbb{P}^n$ as a manifold, and the orbifold structure is given by a $\mathbb{Q}$ -divisor

$$\Delta = \sum_{i=0}^{n+1} \left(1 - \frac{1}{m_i}\right) D_i,$$

where

$$D_i = \{z_i = 0\} \quad \text{for} \quad i = 0, \dots, n, \quad D_{n+1} = \{z_0 + \dots + z_n = 0\},$$

and the $m_0, \dots, m_{n+1}$ are pairwise relatively prime ramification indices. (See Section 4 for precise definitions.) The orbifold first Chern class is

$$c_1(\mathbb{P}^n, \Delta) = (n+1) - \sum_{i=0}^{n+1} \left(1 - \frac{1}{m_i}\right) = \sum_{i=0}^{n+1} \frac{1}{m_i} - 1,$$

where we have identified $H^2(\mathbb{P}^n, \mathbb{Q})$ with $\mathbb{Q}$. Thus $c_1(\mathbb{P}^n, \Delta)$ is positive iff

$$\sum_{i=0}^{n+1} \frac{1}{m_i} - 1 > 0. \tag{1}$$

The existence result [10, Theorem 34] shows that $(\mathbb{P}^n, \Delta)$ has an orbifold Kähler-Einstein metric if in addition the following inequality is also satisfied

$$\sum_{i=0}^{n+1} \frac{1}{m_i} - 1 < \frac{n+1}{n} \min_i \left\{ \frac{1}{m_i} \right\}. \tag{2}$$

This paper started with the observation that one can apply the method of [1] to the *identity* map $(\mathbb{P}^n, \Delta) \rightarrow \mathbb{P}^n$ which is a Galois cover (with trivial

Galois group). On the other hand, over the affine chart $\mathbb{P}^n \setminus \{D_i \cup D_j\}$ the same map can be viewed as having cyclic Galois group of order $\prod_{k \neq i, j} m_k$. This approach improves the bound of [10] by a factor of n , and we obtain

Theorem 1 *Let $D_0, \dots, D_{n+1} \subset \mathbb{P}^n$ be hyperplanes in general position and $m_0, \dots, m_{n+1}$ pairwise relatively prime natural numbers. Assume that*

$$0 < \sum_{i=0}^{n+1} \frac{1}{m_i} - 1 < (n+1) \min_i \left\{ \frac{1}{m_i} \right\}. \quad (3)$$

Then there is an orbifold Kähler-Einstein metric on $(\mathbb{P}^n, \sum_{i=0}^{n+1} (1 - \frac{1}{m_i}) D_i)$.

Set $M = \prod_i m_i$ and $w_i = M/m_i$. As shown in [10] the intersection of the unit sphere with the Brieskorn–Pham singularity

$$L(m_0, \dots, m_{n+1}) := S^{2n+3} \cap \left(\sum_{i=0}^{n+1} z_i^{m_i} = 0 \right) \subset \mathbb{C}^{n+2}$$

is homeomorphic to S^{2n+1} and a Kähler-Einstein metric on the corresponding projective orbifold

$$(X, \Delta_X) := \left(\left(\sum_{i=0}^{n+1} z_i^{m_i} = 0 \right), \sum_{i=0}^{n+1} \left(1 - \frac{1}{m_i} \right) [z_i = 0] \right) \subset \mathbb{P}(w_0, \dots, w_{n+1})$$

lifts to a positive Ricci curvature Einstein metric on $L(m_0, \dots, m_{n+1})$. The weighted projective space $\mathbb{P}(w_0, \dots, w_{n+1})$ is not well formed and it is isomorphic to the ordinary projective space $\mathbb{P}^{n+1}$ by the map

$$(z_0, \dots, z_{n+1}) \mapsto (x_0 = z_0^{m_0}, \dots, x_{n+1} = z_{n+1}^{m_{n+1}}).$$

Under this isomorphism we get that

$$(X, \Delta_X) \cong \left(\left(\sum_{i=0}^{n+1} x_i = 0 \right), \sum_{i=0}^{n+1} \left(1 - \frac{1}{m_i} \right) [x_i = 0] \right) \subset \mathbb{P}^{n+1}.$$

By eliminating the variable x_{n+1} we get that

$$(X, \Delta_X) \cong (\mathbb{P}^n, \Delta).$$

The isometry class of the metric on the sphere determines the complex orbifold $(\mathbb{P}^n, \sum_{i=0}^{n+1} (1 - \frac{1}{m_i}) D_i)$, except possibly when $(\mathbb{P}^n, \sum_{i=0}^{n+1} (1 - \frac{1}{m_i}) D_i)$ has a holomorphic contact structure. The latter can happen only when n is odd;

see [10, Lem.17] for another necessary condition. (Note that $n + 2$ hyperplanes in general position do not have moduli, so the numbers $m_0, \dots, m_{n+1}$ alone determine the complex orbifold.)

Even with the improved bounds, the equations (3) are not easy to satisfy. Still, as in Example 43, we get 12 new Einstein metrics on S^5 corresponding to the ramification indices

$$m_0 = 2, m_1 = 3, m_2 = 5, m_3 \in \{17, 19, 23, 29, 31, 37, 41, 43, 47, 49, 53, 59\},$$

$\geq 10^3$ new Einstein metrics on S^7 , $\geq 10^6$ new Einstein metrics on S^9 ...

The above construction can be varied in many ways. For instance, one can take more than $n + 2$ hyperplanes and quadrics. In all of these cases one gets an improvement by a factor roughly n compared to the bounds in [10], but this gives many new cases only for n large. (As shown by Orlik and Wagreich [21], taking higher degree hypersurfaces for the D_i yields Einstein metrics on various rational homology spheres.)

As another application, we consider singular degree 2 Del Pezzo surfaces. These are all double covers of $\mathbb{P}^2$ ramified along a quartic curve. In the smooth case the existence of Kähler-Einstein metrics was proved by Tian [25]. For singular surfaces we get the following.

Theorem 2 *Let S be a degree 2 Del Pezzo surface with only A_1 or A_2 singularities. Then S has an orbifold Kähler-Einstein metric.*

Anyone well versed in orbifolds, stacks and in the theory of Monge-Ampère equations should have no problem developing the theory of [1] in the orbifold setting. Nonetheless, since the theory of orbifolds has too many “well known” but never proved theorems and not quite correct definitions and proofs, we felt that it makes sense to write down the arguments in some detail.

2 Analytic coverings

Let X and Y be reduced complex spaces. A map $\pi : X \rightarrow Y$ is called *finite* if it is proper and has finite fibres. Since X is locally compact a finite to one map is proper if and only if it is closed. Therefore a map is finite if and only if it is closed and has finite fibres. (By contrast note that $\pi : \mathbb{C} \setminus \{-1\} \rightarrow \{y^2 = x^3 + x^2\} \subset \mathbb{C}^2$ given by $t \mapsto (t^2 - 1, t^3 - t)$ is a closed map of algebraic varieties with finite fibers but π is not proper.)

The fundamental theorem on finite maps (see [15, p. 179]) states that when X and Y are irreducible any finite surjective map $\pi : X \rightarrow Y$ is an *analytic covering*. This means that there is a thin subset $T \subset Y$ such that

- a) $\pi^{-1}(T)$ is thin in X , and
- b) the restriction $\pi^{-1}(Y \setminus T) \rightarrow Y \setminus T$ is locally biholomorphic (étale).

Put $Y_0 = Y \setminus T$ and $X_0 = \pi^{-1}(Y_0)$. Then $\pi : X_0 \rightarrow Y_0$ is a topological covering. We call it a *regular subcover* of π .

We assume that our spaces are irreducible so that “analytic covering” and “finite holomorphic surjection” can be regarded as synonyms.

Another important fact is that an analytic covering $\pi : X \rightarrow Y$ with X and Y normal is an open map (see [15, p. 135]).

Let now $\pi : X \rightarrow Y$ be an analytic covering among connected *normal* complex spaces. Put $Y' = \{y \in Y_{\text{reg}} : \pi^{-1}(y) \subset X_{\text{reg}}\}$ and $X' = \pi^{-1}(Y')$. Then X' and Y' are open sets with complements of codimension at least 2. Now $\pi : X' \rightarrow Y'$ is a finite surjective map between complex manifolds. Pick local coordinates $z_1, \dots, z_n$ on a neighbourhood U of a point in X' and let $w_1, \dots, w_n$ be coordinates around its image in Y' . Let $w_i = \pi_i(z)$ be the local expression of π . The divisors locally defined by the equation

$$\det \left(\frac{\partial \pi_i}{\partial z_j} \right) = 0$$

glue together yielding a well-defined divisor on X' . Since the complement of X' has codimension at least 2, the Remmert-Stein extension theorem (see e.g. [15, p. 181]) ensures that the topological closure of this divisor is a divisor in X , called the *ramification divisor* of π , and denoted by $R = R(\pi)$. It satisfies the Hurwitz formula $K_{Y'} = \pi^* K_{X'} + R$. Write $R = \sum_j r_j R_j$ with R_j distinct prime divisors on X' . The reduced divisor $R_{\text{red}} = \sum_j R_j$ is called the *ramification locus*. By the implicit function theorem $R_{\text{red}} \cap X'$ is the set of points $x \in X'$ such that π is not étale at x , that is the set of critical points of π . Since π is finite, the image $\pi(R_{\text{red}})$ is a divisor on Y , called the *branch divisor* of π .

Consider now the sets $X'' = X' \setminus ((R_{\text{red}})_{\text{sing}} \cup \pi^{-1}(B_{\text{sing}}))$ and $Y' = \pi(X'')$. Both are open and have complements of codimension at least 2 in X and Y respectively. We use this notation often in the sequel. When we want to stress the dependence on π , we write $X''(\pi)$ and $Y''(\pi)$. If $x \in X''$ either $x \notin R_{\text{red}}$ or x belongs to one and only one component R_j . In the first case we say that π is *unramified* at x , in the latter case we say that the *ramification order of π at x is $r_j + 1$* . The ramification order of π at x will

be denoted by $\text{ord}_\pi(x)$. When π is unramified at x , we put $\text{ord}_\pi(x) = 1$. If $D \subset X$ is an irreducible divisor, then there is an open dense subset $D'' \subset D$ such that $\text{ord}_\pi(x)$ does not depend on $x \in D''$. This common value is denoted by $\text{ord}_\pi(D)$ and it is called the *ramification order of π along D* .

We use some basic properties of analytic coverings and maps between them (see, for instance, [6, Lemma 16.1]).

Lemma 3 *Let $x \in X''$. If π is unramified at x , then π is a local biholomorphism at x . If it has ramification order $m > 1$, let R_j be the component of R_{red} passing through x . Then there are local coordinates $z_1, \dots, z_n$ on X'' and $w_1, \dots, w_n$ on Y'' centred at x and $y = \pi(x)$ respectively, such that locally $R_j = \{z_1 = 0\}$, $B = \{w_1 = 0\}$ and $\pi(z_1, \dots, z_n) = (z_1^m, z_2, \dots, z_n)$.*

Since the complement of X'' has codimension 2, R_{red} is the closure of $R_{\text{red}} \cap X''$, that is the closure of the set of points where π has ramification order > 1 .

The next lemma considers the problem of lifting in the simplest case. Denote by $D(r)$ the disc of radius r centred at the origin, by $D^*(r)$ the complement of $\{0\}$ in $D(r)$, and by $P(r_1, \dots, r_n)$ the polydisc centred at the origin with polyradius $(r_1, \dots, r_n)$.

Lemma 4 *Let $P_1 = P(r_1, \dots, r_n)$, $P_2 = P(\rho_1, \dots, \rho_n)$, $Q_1 = P(r_1^{m_1}, r_2, \dots, r_n)$, $Q_2 = P(\rho_1^{m_2}, \rho_2, \dots, \rho_n)$. Set $P_1^* = D^*(r_1) \times P(r_2, \dots, r_n)$ and similarly for P_2^*, Q_1^*, Q_2^* . Let $\pi_i : P_i \rightarrow Q_i$ be the maps $\pi_1(z_1, \dots, z_n) = (z_1^{m_1}, z_2, \dots, z_n)$, $\pi_2(z_1, \dots, z_n) = (z_1^{m_2}, z_2, \dots, z_n)$. Let $f : Q_1 \rightarrow Q_2$ be a holomorphic map such that $f(Q_1^*) \subset Q_2^*$. If $m_2 | m_1$ there are exactly m_2 liftings of f (that is maps $\tilde{f} : P_1 \rightarrow P_2$ such that $\pi_2 \tilde{f} = f \pi_1$). Any local lifting of f defined in a neighbourhood of some point $x \in P_1$ extends to one of these liftings defined on P_1 .*

Lemma 5 *Let $\pi_1 : X_1 \rightarrow Y$ and $\pi_2 : X_2 \rightarrow Y$ be analytic coverings. For $U \subset X_1$ set*

$$\mathfrak{F}(U) = \{\text{holomorphic maps } s : U \rightarrow X_2 \text{ such that } \pi_1 = \pi_2 \circ s\}.$$

Then $\mathfrak{F}$ is a Hausdorff sheaf (of sets) over X_1 . Assume that for any $x_1 \in X_1'', x_2 \in X_2''$ with $\pi_1(x_1) = \pi_2(x_2)$

$$\text{ord}_{\pi_2}(x_2) | \text{ord}_{\pi_1}(x_1).$$

Then the restriction of $\mathfrak{F}$ to $X_1'' \cap \pi_1^{-1}Y_2''(\pi_2)$ is a finite topological covering. In particular, if X_1'' is simply connected, then for every $x_1 \in X_1'' \cap \pi_1^{-1}Y_2''(\pi_2)$ and $x_2 \in X_2''$ such that $\pi_1(x_1) = \pi_2(x_2)$ there is an analytic map $f : X_1'' \rightarrow X_2$ such that $f(x_1) = x_2$ and $\pi_1 = \pi_2 \circ f$.

In fact, the above f extends to X_1 by the following immediate consequence of the Riemann Extension Theorem (see e.g. [15, p.144])

Lemma 6 *Let $\pi_1 : X_1 \rightarrow Y$ and $\pi_2 : X_2 \rightarrow Y$ be analytic coverings, X_1 normal and $T \subset X_1$ a thin set. Let $f^0 : X_1 \setminus T \rightarrow X_2$ be an analytic map such that $\pi_1 = \pi_2 \circ f^0$. Then f^0 extends to $f : X_1 \rightarrow X_2$ such that $\pi_1 = \pi_2 \circ f$.*

3 The Galois group of coverings

Let $f : X \rightarrow Y$ be an analytic covering of normal complex spaces. Put $\text{Gal}(\pi) = \{f \in \text{Aut}(X) : \pi \circ f = \pi\}$. $\text{Gal}(\pi)$ is a finite subgroup of $\text{Aut}(X)$. In fact fix $x \in X''$, $y = \pi(x)$, and let V be a neighbourhood of y in Y such that $\pi^{-1}(V) = \bigcup_{i=1}^k U_i$ with $\pi : U_i \rightarrow V$ a biholomorphism and $x \in U_1$. Then the stabiliser $\text{Gal}(\pi)_x$ is a subgroup of finite index in $\text{Gal}(\pi)$. Moreover any $f \in \text{Gal}(\pi)_x$ maps U_1 to itself. Since $\pi|_{U_1}$ is injective, the restriction of f to U_1 is the identity. By the connectedness of X , $f = \text{id}_X$, so $\text{Gal}(\pi)_x = \{1\}$ and $\text{Gal}(\pi)$ is finite.

Since π is $\text{Gal}(\pi)$ -invariant, the $\text{Gal}(\pi)$ -orbit of $x \in X$ is contained in $\pi^{-1}(\pi(x))$. We say that an analytic covering $\pi : X \rightarrow Y$ is *Galois* if the converse holds, that is two points of X lie on the same fibre of π only if they belong to the same $\text{Gal}(\pi)$ -orbit.

The branching divisor of a Galois cover can be described also in the following way. Given a prime divisor D in X , set $\Gamma(D) = \{\gamma \in \Gamma : D \subset \text{Fix}(\gamma)\}$. For each prime divisor D the image $\pi(D)$ is a prime divisor in Y . The prime divisors for which $\Gamma(D) \neq 0$ are exactly the R_j . Set $B_j = \pi(R_j)$. In general different R_j 's can have the same image. Assume that $\{B_i\}_{i \in I}$ is the set of all images of the R_j 's (that is $B_i \neq B_k$ if $i \neq k$). Then

$$B(\pi) = \sum_{i \in I} \left(1 - \frac{1}{|\Gamma(R_i)|}\right) B_i. \quad (4)$$

Lemma 7 *Let X and Y be normal complex spaces, $\pi : X \rightarrow Y$ an analytic covering and $Y_0 \subset Y$ an open subset with thin complement. Put $X_0 = \pi^{-1}(Y_0)$ and $\pi_0 = \pi|_{X_0} : X_0 \rightarrow Y_0$. Then the elements of $\text{Gal}(\pi_0)$ extend to elements of $\text{Gal}(\pi)$, and if π_0 is Galois, then π is Galois too.*

Proof. The first part follows from Lemma (6). For the second part, let $x, x' \in X$ be such that $\pi(x) = \pi(x') = y$. If $y \in Y_0$ there is some $g \in \text{Gal}(\pi_0)$ such that $g.x = x'$. Since we have just proved that $\text{Gal}(\pi_0) = \text{Gal}(\pi)$ the Galois condition is satisfied for these points. If instead $y \in Y \setminus Y_0$, choose

neighbourhoods U_i and V as above. Assume $x = x_1 \in U_1$ and $x' = x_2 \in U_2$. Let $\{z_n\}$ be a sequence of points in $X_0 \cap U_1$ converging to x . Then $y_n = \pi(z_n)$ converge to y . Since π is open, $\pi(U_2) = V$. Therefore there are points $z'_n \in U_2 \cap X_0$ such that $\pi(z'_n) = y_n$. By the Galois condition on X_0 , there are $g_n \in \text{Gal}(\pi)$ such that $z'_n = g_n \cdot z_n$. As $\text{Gal}(\pi)$ is finite, we can extract a subsequence with $g_n \equiv g$. Since $\lim z'_n = x_2$ as $\pi^{-1}(y) \cap U_2 = \{x_2\}$, we get $x_2 = g \cdot x_1$. $\square$

If $\pi : X \rightarrow Y$ is a Galois covering, then $\text{Gal}(\pi)$ acts freely on any regular subcover X_0 . Therefore if $x, x' \in X_0$ and $\pi(x) = \pi(x')$, then there is a unique $g \in \text{Gal}(\pi)$ such that $g \cdot x = x'$. In particular the cardinality of $\text{Gal}(\pi)$ equals that of the generic fibre. This condition is also sufficient: π is Galois iff $|\text{Gal}(\pi)|$ equals the cardinality of the general fibre iff $\text{Gal}(\pi)$ is transitive on the general fibre.

For later reference we state the following simple lemma.

Lemma 8 *Let X, Y and Z be irreducible complex spaces, and $f : X \rightarrow Z$, $g : Y \rightarrow Z$, $h : X \rightarrow Y$ analytic coverings such that $gh = f$. If f is Galois, then h is Galois too.*

Proof. Thanks to Lemma 7 it is enough to consider the unramified case. Fix $x \in X$ and put $y = h(x), z = f(x) = g(y)$. We need to show that $h_*\pi_1(X, x)$ is a normal subgroup of $\pi_1(Y, y)$. Since $g_* : \pi_1(Y, y) \rightarrow \pi_1(Z, z)$ is injective it is enough to check that $g_*h_*\pi_1(X, x)$ is a normal subgroup of $g_*\pi_1(Y, y)$. But f being Galois $f_*\pi_1(X, x) = g_*h_*\pi_1(X, x)$ is normal in $\pi_1(Z, z)$, hence a fortiori in $g_*\pi_1(Y, y)$. $\square$

For a general analytic covering $\pi : X \rightarrow Y$ it is not possible to assign multiplicity to the branching divisor in any reasonable way. In fact, different points in the preimage of a point $y \in B$ have different branching orders. A typical example is $X = \{z^3 - 3yz + 2x = 0\} \subset \mathbb{C}^3$ projecting on $\mathbb{C}_{x,y}^2$. Even shrinking the domain around the origin, one cannot separate the branches with different orders.

On the other hand, when the covering is Galois, for any $y \in Y''$ all points in $\pi^{-1}(y)$ have the same branching order. Therefore we can assign multiplicities to the branch divisor according to the following rule. Let $y \in Y'' \cap B$ and let x be any point in $\pi^{-1}(y)$. Then we define the multiplicity of B in y to be $1 - 1/\text{ord}_\pi(x)$. We still denote by B the $\mathbb{Q}$ -divisor given by the branching locus provided with these multiplicities. Note that with this convention $R = \pi^*B$, that is, the ramification divisor is the pull back of the branch divisor.

4 Orbifolds as pairs

As in [10], we look at orbifolds as a particular type of log pairs. (X, Δ) is a log pair if X is a normal algebraic variety (or a normal complex space) and $\Delta = \sum_i d_i D_i$ is an effective $\mathbb{Q}$ -divisor where the D_i are distinct, irreducible divisors and $d_i \in \mathbb{Q}$. The number d_i is called the multiplicity of Δ along D_i , it is denoted by $\text{mult}_{D_i} \Delta$. We set $\text{mult}_D \Delta = 0$ for every other irreducible divisor $D \neq D_i \forall i$.

Let $X''(\Delta)$ (or simply X'') be the complement of $X_{\text{sing}} \cup \Delta_{\text{sing}}$. For $x \in X''$ the multiplicity of Δ at x is a well defined rational number. For orbifolds, we need to consider only pairs (X, Δ) such that Δ has the form

$$\Delta = \sum_i (1 - \frac{1}{m_i}) D_i,$$

where the D_i are prime divisors and $m_i \in \mathbb{N}$. If (X, Δ) is such a pair then for any divisor $D \subset X$ we put

$$\text{ord}_\Delta(D) = \frac{1}{1 - \text{mult}_D \Delta}.$$

The assumption on the multiplicities of Δ amounts to saying that the order is always a nonnegative integer.

Definition 9 *An orbifold chart on X compatible with Δ is a Galois covering $\varphi : U \rightarrow \varphi(U) \subset X$ such that*

1. *U is a domain in $\mathbb{C}^n$ and $\varphi(U)$ is open in X ;*
2. *the branch locus of φ is $\Delta_{\text{red}} \cap \varphi(U)$;*
3. *for any $x \in U''(\varphi)$ such that $\varphi(x) \in D_i$, $\text{ord}_\varphi(x) = m_i$.*

Conditions (2) and (3) are equivalent to

$$B(\varphi) = \Delta \cap \varphi(U). \tag{5}$$

Definition 10 *An orbifold is a log pair (X, Δ) such that X is covered by orbifold charts compatible with Δ .*

(For a slightly more general approach, see [13, §14].)

Let X be a normal complex space and $\pi : U \rightarrow X$ a Galois cover where U is a smooth. As discussed earlier, the branch divisor $B(\pi)$ of π is defined and we get a log pair $(X, B(\pi))$. If U is simply connected, (which we can always assume by shrinking U suitably) then by Lemma 5 the log pair $(X, B(\pi))$ determines $\pi : U \rightarrow X$ up to biholomorphisms. Thus we recover the classical definition of orbifolds (as in [4] for example).

Example 11 Let X be a complex manifold and $D = \sum_{i \in I} D_i$ a divisor with local normal crossing. By this we mean that for any point $x \in X$ there is a holomorphic coordinate system $(V, z_1, \dots, z_n)$ such that $D \cap V = \{z \in V : z_1 \cdots z_k = 0\}$. If $D_i \cap V \neq \emptyset$ then $D_i \cap V$ is the union of some of the hypersurfaces $\{z_j = 0\}$. (D is said to be a divisor with global normal crossing if, in addition, each D_i is smooth.) For any $i \in I$, fix an integer $m_i > 1$ and put $\Delta = \sum_i (1 - 1/m_i) D_i$. We claim that (X, Δ) is an orbifold. Indeed, fix a coordinate system as above and put $m'_j = m_i$ if $\{z_j = 0\} \subset D_i \cap V$. Set

$$\varphi : U \rightarrow V, \quad \pi(x_1, \dots, x_n) = (x_1^{m'_1}, \dots, x_k^{m'_k}, x_{k+1}, \dots, x_n). \quad (6)$$

Then (U, φ) is orbifold chart on X compatible with Δ and so (X, Δ) is an orbifold.

In the same way, the usual definition of orbifold map is equivalent to the following one.

Definition 12 A finite holomorphic map $f : X \rightarrow Y$ is an orbifold map $f : (X, \Delta_X) \rightarrow (Y, \Delta_Y)$ if

$$\text{ord}_{\Delta_Y}(f(D)) \mid \text{ord}_{\Delta_X}(D) \cdot \text{ord}_f D \quad (7)$$

for every divisor $D \subset X$.

An orbifold automorphism is an orbifold map that is invertible with inverse an orbifold map. The group of automorphisms of (X, Δ) is denoted by $\text{Aut}(X, \Delta)$.

Definition 13 An orbifold Galois covering $f : (X, \Delta_X) \rightarrow (Y, \Delta_Y)$ is an orbifold map such that $f : X \rightarrow Y$ is a Galois analytic cover and $\text{Gal}(f) \subset \text{Aut}(X, \Delta_X)$.

By the *degree* of an orbifold Galois cover we mean its *degree* as an analytic cover.

Lemma 14 Let $f : (X, \Delta_X) \rightarrow (Y, \Delta_Y)$ be an orbifold map. Then given $x \in X$ and $y = f(x) \in Y$ there are orbifold charts (U, φ) and (V, ψ) around x and y respectively such that f has a lifting $\tilde{f} : U \rightarrow V$. If, in addition, $f : X \rightarrow Y$ is a Galois covering then $\tilde{f} : U \rightarrow V$ is also a Galois covering.

Proof. Choose the chart (U, φ) such that U is simply connected and $f(\varphi(U)) \subset \psi(V)$. If $D \subset U$ is any divisor then

$$\text{ord}_{f \circ \varphi} D = \text{ord}_f \varphi(D) \cdot \text{ord}_\varphi D = \text{ord}_f \varphi(D) \cdot \text{ord}_{\Delta_X} \varphi(D).$$

By the definition of orbifold maps,

$$\text{ord}_{\Delta_Y}(f \circ \varphi)(D) \mid \text{ord}_f \varphi(D) \cdot \text{ord}_{\Delta_X} \varphi(D),$$

hence we conclude that $\text{ord}_{\Delta_Y}(f \circ \varphi)(D)$ divides $\text{ord}_{f \circ \varphi} D$. Thus the assumption of Lemma 5 is satisfied and so $f \circ \varphi$ lifts to $\tilde{f} : U \rightarrow V$. Assume next that $f : X \rightarrow Y$ is a Galois covering and pick $u_1, u_2 \in U$ such that $\tilde{f}(u_1) = \tilde{f}(u_2)$. Then $f(\varphi(u_1)) = f(\varphi(u_2))$ hence there is a Galois automorphism σ of f such that $\varphi(u_1) = \sigma(\varphi(u_2))$. Applying Lemma 5 to $\varphi : U \rightarrow X$ and $\sigma \circ \varphi : U \rightarrow X$ we conclude that σ lifts to a biholomorphism $\tilde{\sigma}$ of U such that $\varphi(u_1) = \varphi(\tilde{\sigma}(u_2))$. Since $\varphi : U \rightarrow X$ is Galois, there is a biholomorphism ρ of U such that $u_1 = \rho(\tilde{\sigma}(u_2))$. This shows that in the commutative diagram

$$\begin{array}{ccc} U & \xrightarrow{\tilde{f}} & V \\ \downarrow \varphi & & \downarrow \psi \\ \varphi(U) & \xrightarrow{f} & \psi(V). \end{array} \quad (8)$$

the composite $f \circ \varphi$ is Galois. But $f \circ \varphi = \psi \tilde{f}$ and by Lemma 8 $\tilde{f}$ is a Galois cover. $\square$

Example 15 Let (X, Δ) be any orbifold, and let $(X, 0)$ denote the orbifold structure on X with trivial branching divisor. It is a nontrivial result that $(X, 0)$ is an orbifold, that is, X has quotient singularities (see [22]). (We use mainly the case when X is smooth, and then the orbifold charts of $(X, 0)$ are simply the manifold charts of X .)

The identity map $\text{id}_X : (X, \Delta) \rightarrow (X, 0)$ is trivially an orbifold Galois covering. In fact it is both an orbifold map and a Galois analytic cover, and $\text{Gal}(\text{id}_X) = \{\text{id}_X\} \subset \text{Aut}(X, \Delta)$.

If $f : (X, \Delta) \rightarrow (Y, \Delta_Y)$ is an orbifold Galois covering the *orbifold ramification divisor* of f is defined as

$$R^{\text{orb}}(\Delta_X, \Delta_Y, f) = R(f) + \Delta_X - f^* \Delta_Y.$$

With this definition the logarithmic ramification formula

$$K_X + \Delta_X = f^*(K_Y + \Delta_Y) + R^{orb}(\Delta_X, \Delta_Y, f)$$

is automatically satisfied. To understand the geometric meaning of R^{orb} it is useful to look at the open set

$$X''(\Delta_X, \Delta_Y, f) = X_{\text{reg}} \cap f^{-1}(Y_{\text{reg}} \setminus (\Delta_Y \cup B(f))_{\text{sing}}) \setminus (\Delta_X \cup R(f))_{\text{sing}}.$$

This means that $x \in X'' = X''(\Delta_X, \Delta_Y, f)$ if (a) X is smooth at x , (b) Y is smooth at $y = f(x)$, (c) x belongs to at most one component D of $\Delta_X + R(f)$ and in this case x is a smooth point of D , (d) y belongs to at most one component D' of $\Delta_Y + B(f)$ and in this case it is a smooth point of D' . As usual the complement of this set has codimension 2. Let D be any smooth divisor passing through x and D' a smooth component passing through y . Assume first that f is unbranched at x and that locally $\Delta_X = (1 - 1/p)D$ and $\Delta_Y = (1 - 1/q)D'$. Then there is a local diagram like (8), with $p = \deg \varphi$ and $q = \deg \psi$. Put $k = \deg \tilde{f}$. Since f is unbranched we can assume that its restriction to $\varphi(U)$ is a biholomorphism onto $\psi(V)$. Therefore $p = qk$. If $p = 1$, then $q = k = 1$, and as expected $\text{mult}_x R^{orb} = 0$. If $p > 1$, then necessarily $D' = f(D)$ because of (7) and $f^*D' = D$, since f is étale. Therefore $R^{orb} = (1/q - 1/p)D = (k - 1)/p \cdot D$. If instead $\text{ord}_x(f) = m > 1$, then again $D' = f(D)$, $R(f) = (m - 1)D$, $f^*D' = mD$, $pm = qk$ and $R^{orb} = (m/q - 1/p)D = (k - 1)/p \cdot D$ once more. Roughly the orbifold ramification divisor is the ramification of the lifting $\tilde{f}$ divided the degree of the local chart φ .

Let (X, Δ) be an orbifold and $\Gamma \subset \text{Aut}(X, \Delta)$ a finite subgroup. We want to define a quotient orbifold (Y, Δ') . By Cartan's lemma [11] $Y = X/\Gamma$ is a normal analytic space and the canonical projection $\pi : X \rightarrow Y$ is an analytic covering. The support of the branch divisor Δ' is defined to be $\pi(\Delta) \cup B(\pi)$, while the multiplicities are specified as follows. Let D be an irreducible component of $\pi(\Delta) \cup B(\pi)$. If D is a component of $\pi(\Delta)$ and not of $B(\pi)$, then we assign to D the multiplicity $\text{mult}_x(\Delta)$, where x is any point in $X''(\Delta)$ such that $\pi(x) \in D$ is a smooth point of $\pi(\Delta) \cup B(\pi)$. If D is a component of $B(\pi)$ and not of $\pi(\Delta)$ then we assign to D the same multiplicity it has as a component of $B(\pi)$, that is $1 - 1/\text{ord}_\pi(x)$ for any $x \in X''(\pi)$ such that $\pi(x) \in D$ is a smooth point of $\pi(\Delta) \cup B(\pi)$. Finally, if D is a common component of $\pi(\Delta)$ and $B(\pi)$ then we assign to it the multiplicity

$$1 - \frac{1 - \text{mult}_x \Delta}{\text{ord}_\pi(x)}$$

for any $x \in X''(\Delta) \cap X''(\pi)$ such that $\pi(x) \in D$ is a smooth point of $\pi(\Delta) \cup B(\pi)$.

Proposition 16 *Let (X, Δ) be an orbifold, and $\Gamma \subset \text{Aut}(X, \Delta)$ a finite subgroup. Let $Y = X/\Gamma$ be the quotient analytic space, and Δ' the $\mathbb{Q}$ -divisor defined above. Then (Y, Δ') is an orbifold and the canonical projection*

$$\pi : (X, \Delta_X) \longrightarrow (Y, \Delta') \quad (9)$$

is an orbifold Galois covering.

Proof. We need to show that Y is covered by orbifold charts compatible with Δ' . Fix $y \in Y$, $x \in \pi^{-1}(y)$ and let $\varphi : U \rightarrow \varphi(U)$ be an orbifold chart with $x \in \varphi(U)$. If the stabiliser Γ_x is trivial we can assume that $\gamma\varphi(U) \cap \varphi(U) = \emptyset$ for any $\gamma \neq e$. Then $\pi : \varphi(U) \rightarrow Y$ is a biholomorphism onto its image. Put $\psi = \pi\varphi : U \rightarrow Y$. We claim that ψ is an orbifold chart on Y compatible with Δ' . In fact ψ is Galois since π is a biholomorphism on $\varphi(U)$, and $\pi^*B(\psi) = B(\varphi) = \Delta \cap \varphi(U)$. On the other hand $B(\pi) \cap \psi(U) = \emptyset$ since $\pi : \varphi(U) \rightarrow \psi(U)$ is biholomorphic. Therefore on $\psi(U)$ the divisor Δ' coincides with $B(\psi)$. This proves that $\psi : U \rightarrow Y$ is an orbifold chart. If $\Gamma_y \neq \{e\}$ take a chart $\varphi : U \rightarrow \varphi(U) \subset X$ such that $\varphi(U)$ be a Γ_x -invariant neighbourhood of x . Lemma 14 ensures that also in this case $\psi = \pi\varphi : U \rightarrow \psi(U) \cong \varphi(U)/\Gamma_x$ is a Galois covering. It is easy to verify that $B(\psi) = \Delta'$ on $\psi(U)$. Finally that π is an orbifold Galois covering is clear: a lifting of $\pi : \varphi(U) \rightarrow \psi(U)$ is given by the identity map $U \rightarrow U$ which is trivially Galois. $\square$

5 Basic estimates for orbifold Kähler-Einstein metrics

In this section we collect the orbifold versions of some fundamental results due to Aubin, Bando-Mabuchi and Tian, that are needed in the existence criteria in the next section. Most of the proofs are the same as in the case of a manifold and we just give appropriate references. For the basic definitions of differential geometry on orbifolds see [4], [3], [9] and [7]. Some information on Sobolev spaces and Laplace operators on orbifolds can be found e.g. in [12].

Remark 17 *Note that if X is a complex manifold and Δ is a non trivial branching divisor, then smoothness in the orbifold sense is rather different from ordinary smoothness. For example, $f(z) = |z|$ is not smooth*

in the ordinary sense, but it belongs to $C^\infty(\mathbb{C}, \Delta)$, where Δ is the divisor concentrated at the origin with multiplicity $1/2$. In fact the inclusions $C^\infty(X) \subsetneq C^\infty(X, \Delta)$ and $\wedge^k(X) \subsetneq \wedge^k(X, \Delta)$ are in general strict.

Definition 18 A Fano orbifold is a compact complex orbifold (X, Δ) such that $-(K_X + \Delta)$ is ample.

By the Baily-Kodaira imbedding theorem [3] this is equivalent to the fact that $c_1(X, \Delta)$ contains an orbifold Kähler metric.

The following is the orbifold analogue of Bonnet-Myers Theorem. It follows, for example, from the Bishop volume comparison Theorem for orbifolds, see [7, Prop. 20, Cor. 21].

Theorem 19 Let X be an m -dimensional orbifold and g a Riemannian orbifold metric on X with $\text{Ric}(g) \geq \varepsilon(m-1)g$ for some $\varepsilon > 0$. Then $\text{diam}(X, g) \leq \pi/\sqrt{\varepsilon}$.

Theorem 20 ([20, Theorem B]) Let (X, g) be a Riemannian orbifold of dimension $m > 2$ with $\text{Ric}(g) \geq -(m-1)\varepsilon^2 g$ for some $\varepsilon \geq 0$. Then there is a constant $C > 0$ depending only on m and $\varepsilon \cdot \text{diam}(X, g)$ such that

$$\|\nabla u\|_{L^2} \geq C \frac{\text{vol}(X, g)^{1/m}}{\text{diam}(X, g)} \|u\|_{L^{2m/(m-2)}} \quad (10)$$

for any $u \in W^{1,2}(X)$ with $\int_X u \, d\text{vol}_g = 0$.

Combining the last two theorems one gets the following uniform Sobolev embedding.

Corollary 21 Let (X, Δ) be an n -dimensional Fano orbifold. For any $\varepsilon > 0$ there is a constant $C = C(\varepsilon) > 0$ such that for any metric ω in the class $2\pi c_1(X, \Delta)$ with $\text{Ric}(\omega) \geq \varepsilon\omega$ and any $u \in W^{1,2}(X, \Delta)$

$$\|u\|_{L^{2n/(n-1)}} \leq C \|u\|_{W^{1,2}}^2. \quad (11)$$

If (X, Δ) is a Kähler orbifold, $\omega \in \wedge^{1,1}(X, \Delta)$ is a closed smooth form and $\varphi \in C^\infty(X, \Delta)$, put $\omega_\varphi = \omega + i\partial\bar{\partial}\varphi$. We write $\omega_\varphi > 0$ to mean that it is a Kähler metric. If ω is such that

$$\langle [\omega]^n, [X] \rangle = \int_X \omega^n > 0$$

and $\varphi \in C^\infty(X, \Delta)$ put

$$I_\omega(\varphi) = \frac{1}{\langle [\omega]^n, [X] \rangle} \int \varphi(\omega^n - \omega_\varphi^n) \quad (12)$$

$$J_\omega(\varphi) = \int_0^1 \frac{I_\omega(s\varphi)}{s} ds \quad (13)$$

$$F_\omega^0(\varphi) = J_\omega(\varphi) - \frac{1}{\langle [\omega]^n, [X] \rangle} \int \varphi \omega^n. \quad (14)$$

Lemma 22

$$J_\omega(\varphi) = \frac{1}{\langle [\omega]^n, [X] \rangle} \sum_{k=0}^{n-1} \frac{k+1}{n+1} \int_M i \partial \varphi \wedge \bar{\partial} \varphi \wedge \omega^k \wedge \omega_\varphi^{n-k-1} \quad (15)$$

$$I_\omega(\varphi) - J_\omega(\varphi) = \frac{1}{\langle [\omega]^n, [X] \rangle} \sum_{k=0}^{n-1} \frac{n-k}{n+1} \int_X i \partial \varphi \wedge \bar{\partial} \varphi \wedge \omega^k \wedge \omega_\varphi^{n-k-1}. \quad (16)$$

If $\omega > 0$ and $\omega_\varphi > 0$, then $I_\omega(\varphi)$, $J_\omega(\varphi)$ and $I_\omega(\varphi) - J_\omega(\varphi)$ are nonnegative and vanish only if φ is constant. Moreover $J_\omega \leq I_\omega \leq (n+1)J_\omega$.

For (15) see [24, Lemma 2.2] or [1, Lemma 2.1]. For (16) expand $\omega^n - \omega_\varphi^n$. The last statements follow diagonalising simultaneously ω and $i \partial \bar{\partial} \varphi$. $\square$

Lemma 23 *If λ is a positive constant then*

$$F_{\lambda\omega}^0(\lambda\varphi) = \lambda F_\omega^0(\varphi). \quad (17)$$

Let ω_0 be a closed $(1,1)$ -form such that $\langle [\omega_0]^n, [X] \rangle > 0$. Given $\varphi_{01}, \varphi_{12} \in C^\infty(X, \Delta)$ put $\omega_1 = \omega_0 + i \partial \bar{\partial} \varphi_{01}$, $\varphi_{02} = \varphi_{01} + \varphi_{12}$. Then

$$F_{\omega_0}^0(\varphi_{02}) = F_{\omega_0}^0(\varphi_{01}) + F_{\omega_1}^0(\varphi_{12}). \quad (18)$$

(Same proof as in [27, pp. 60f].)

Lemma 24 ([27, p. 59]) *If φ_t is a differentiable family of smooth functions on (X, Δ) then*

$$\frac{d}{dt} J_\omega(\varphi_t) = \frac{1}{\langle [\omega]^n, [X] \rangle} \int_X \dot{\varphi}_t(\omega^n - \omega_t^n) \quad (19)$$

$$\frac{d}{dt} F_\omega^0(\varphi_t) = -\frac{1}{\langle [\omega]^n, [X] \rangle} \int_X \dot{\varphi}_t \omega_t^n \quad (20)$$

Assume now that ω is a Kähler orbifold metric in the canonical class, that is $\omega \in 2\pi c_1(X, \Delta)$. Let $f = f(\omega) \in C^\infty(X, \Delta)$ be the unique function such that

$$\text{Ric}(\omega) - \omega = i \partial \bar{\partial} f(\omega) \quad \int_X e^{f(\omega)} = \int_X \omega^n. \quad (21)$$

Put $V = \langle [\omega]^n, [X] \rangle = n! \text{vol}(X)$ and define $A_\omega, F_\omega : C^\infty(X, \Delta) \rightarrow \mathbb{R}$ by

$$A_\omega(\varphi) = \log \left[\frac{1}{V} \int_X e^{f(\omega) - \varphi} \omega^n \right] \quad F_\omega(\varphi) = F_\omega^0(\varphi) - A_\omega(\varphi). \quad (22)$$

Using the notation of Lemma 23 if ω_0, ω_1 and ω_2 are Kähler metrics, then

$$F_{\omega_0}(\varphi_{02}) = F_{\omega_0}(\varphi_{01}) + F_{\omega_1}(\varphi_{12}). \quad (23)$$

For $G \subset \text{Aut}(X, \Delta)$ a subgroup of isometries of (X, Δ, ω) put

$$P_G(X, \Delta, \omega) = \{\varphi \in C^\infty(X, \Delta) : \omega_\varphi > 0, \text{ and } \varphi \text{ is } G\text{-invariant}\}. \quad (24)$$

If $G = \{1\}$ we simply write $P(X, \Delta, \omega)$.

In order to construct a Kähler-Einstein metric on (X, Δ) the continuity method is applied: fix a Kähler metric ω in the canonical class and consider the well-known equations

$$(\omega + i \partial \bar{\partial} \varphi_t)^n = e^{f - t\varphi_t} \omega^n \quad (*)_t$$

for a smooth family of functions in $C^\infty(X, \Delta)$. Yau's estimates hold for orbifold metrics, and in particular the Calabi conjecture is true, which implies that $(*)_0$ admits a unique solution. Denote by Δ the negative definite $\bar{\partial}$ -Laplacian on functions (that is $\Delta = -\bar{\partial}^* \bar{\partial}$) and by $-\lambda_j$ its eigenvalues.

Lemma 25 ([2, Theorem 4.20 p. 116]) *Let ω be a Kähler metric on the compact orbifold (X, Δ) . If $\text{Ric}(\omega) \geq \varepsilon > 0$, then $\lambda_1 \geq 1$.*

It follows that the times t for which $(*)_t$ is solvable form an open subset $S \subset [0, 1]$ and that solutions φ_t are smooth in t , see [27, pp. 63-66]. Given a C^0 -estimate for the solutions, Yau's estimates ensure that S is closed, thus yielding the solution up to $t = 1$, which is a Kähler-Einstein metric.

Proposition 26 *Let φ_t be a solution to $(*)_t$ for $t \in [0, T_0)$. Then $I_\omega(\varphi_t) - J_\omega(\varphi_t)$ is nondecreasing and $F_\omega^0(\varphi_t) \leq 0$.*

Proof. Differentiating $(*)_t$ with respect to t one gets

$$(\Delta_t + t)\dot{\varphi}_t = -\varphi_t. \quad (25)$$

Therefore

$$\begin{aligned} \frac{d}{dt} (I_\omega(\varphi_t) - J_\omega(\varphi_t)) &= \\ &= \frac{1}{V} \int_X \varphi_t(\varphi_t + t\dot{\varphi}_t)\omega_t^n = (1-t^2) \frac{1}{V} \int_X \varphi_t^2 \omega_t^n + \frac{1}{V} \int_X |\bar{\partial}\dot{\varphi}_t|^2 \omega_t. \end{aligned}$$

This gives the first result. For the second use (20) and (25):

$$\begin{aligned} \frac{d}{dt} tF_\omega^0(\varphi_t) &= F_\omega^0(\varphi_t) - \frac{t}{V} \int_X \dot{\varphi}_t \omega_t^n = \\ &= F_\omega^0(\varphi_t) + \frac{1}{V} \int_X (\Delta_t \dot{\varphi}_t + \varphi_t) \omega_t^n = J_\omega(\varphi_t). \end{aligned}$$

Since $J_\omega \geq 0$ the result follows. $\square$

The following estimates depend on the uniform Sobolev embedding (Lemma 21) and their proof uses Moser iteration.

Theorem 27 ([27, p. 67ff]) *If φ_t is a family of solutions to $(*)_t$ on the time interval $[0, T_0)$, then there is a constant $C = C(T_0) > 0$ such that for any $t < T_0$*

$$\|\varphi_t\|_\infty \leq C(1 + J_\omega(\varphi_t)) \quad (26)$$

$$0 \leq -\inf_X \varphi_t \leq C \left(\frac{1}{V} \int_X (-\varphi_t) \omega_t^n + C \right) \quad (27)$$

$$F_\omega(\varphi_t) \leq -A_\omega(\varphi_t) \leq C(1-t) \leq C. \quad (28)$$

Lemma 28 ([5, §6]) *Let (X, Δ) be a Fano orbifold, ω_{KE} a Kähler-Einstein metric and ω a metric in the canonical class. Then there is $g \in \text{Aut}(X, \Delta)$ such that $\omega = g^* \omega_{KE} + i\partial\bar{\partial}\psi$ with ψ orthogonal to $\ker(\Delta_{g^* \omega_{KE}} + 1)$ in $L^2(X, \omega_{KE}^n)$.*

Proposition 29 ([26, Prop. 5.3]) *Let (X, Δ) be a Fano orbifold and ω_{KE} a Kähler-Einstein metric in the canonical class. If $\omega = \omega_{KE} + i\partial\bar{\partial}\psi$ is a Kähler metric, with $\psi \perp \ker(\Delta_{KE} + 1)$ and $\int_X e^{-\psi} \omega_{KE}^n = 0$, there is a solution $\{\varphi_t\}_{t \in [0,1]}$ of $(*)_t$ with $\varphi_0 = 0$ and $\varphi_1 = -\psi$.*

Theorem 30 (Ding-Tian) *If a Fano orbifold (X, Δ) admits a Kähler-Einstein metric ω_{KE} , then F_ω is bounded from below on $P(X, \Delta, \omega)$ for any ω in the canonical class.*

Proof. Thanks to (23) it is enough to bound $F_{\omega_{KE}}$. Given $\varphi \in P(X, \Delta, \omega_{KE})$ put $\omega = \omega_{KE} + i\partial\bar{\partial}\varphi$ and let g and ψ be as in Lemma 28. Using again (23) it is enough to bound $F_{g^*\omega_{KE}}(\psi)$. Take a path as in Lemma 29. Thanks to Proposition 26 $F_{g^*\omega_{KE}}(\psi) = -F_\omega(-\psi) = -F_\omega(\varphi_1) = F_\omega^0(\varphi_1) \geq 0$. $\square$

Remark 31 *These estimates are enough to prove one half of Tian's fundamental theorem, namely that properness of F_ω implies the existence of a Kähler-Einstein metric (see [27, p. 63]).*

The following normalisation of potentials is useful:

$$Q_G(X, \Delta, \omega) = \{\varphi \in P_G(X, \Delta, \omega) : A_\omega(\varphi) = 0\}. \quad (29)$$

For any $\varphi \in P_G(X, \Delta, \omega)$, $\varphi + A_\omega(\varphi) \in Q_G(X, \Delta, \omega)$.

Proposition 32 *Let (X, Δ) be a Fano orbifold, $\omega \in 2\pi c_1(X, \Delta)$ a Kähler metric and G a compact group of isometries of (X, Δ, ω) . If there are constants $C_1, C_2 > 0$ such that*

$$F_\omega(\varphi) \geq C_1 \sup_X \varphi - C_2 \quad (30)$$

for any $\varphi \in Q_G(X, \Delta, \omega)$, then (X, Δ) admits a Kähler-Einstein metric.

Proof. Let φ_t be a solution of $(*)_t$ on $[0, T_0)$. Since $\varphi_t + A_\omega(\varphi_t) \in Q_G(X, \Delta, \omega)$

$$\begin{aligned} F_\omega(\varphi_t) &= F_\omega(\varphi_t + A_\omega(\varphi_t)) \geq \\ &\geq C_1 \sup_X (\varphi_t + A_\omega(\varphi_t)) - C_2 = C_1 \sup_X \varphi_t + C_1 A_\omega(\varphi_t) - C_2 \end{aligned} \quad (31)$$

Using (28)

$$C_1 \sup_X \varphi_t \leq F_\omega(\varphi_t) - C_1 A_\omega(\varphi_t) + C_2 \leq C_3 + C_2 + C_1 C_3.$$

Hence $\sup_X \varphi_t$ is uniformly bounded. But $F^0(\varphi_t) \leq 0$, so $J_\omega(\varphi_t) \leq F_\omega^0(\varphi_t) + \sup_X \varphi_t$ is bounded and (26) yields the required bound of the C^0 norm. $\square$

Lemma 33 ([1, Lemma 2.3]) *Let (X, Δ) be a Fano orbifold, and $\omega \in 2\pi c_1(M)$ a Kähler metric. Then for any $\beta > 0$ there are constants $C_1, C_2 > 0$ such that for any $\varphi \in Q(X, \Delta, \omega)$*

$$\log \left[\frac{1}{V} \int_X e^{-(1+\beta)\varphi} \omega^n \right] \geq C_1 \sup_X \varphi - C_2. \quad (32)$$

Corollary 34 *If there are constants $C_1, C_2 > 0$ and $\beta > 0$ such that*

$$F_\omega(\varphi) \geq C_1 \log \left[\frac{1}{V} \int_X e^{-(1+\beta)\varphi} \omega^n \right] - C_2 \quad (33)$$

for any $\varphi \in Q_G(X, \Delta, \omega)$, then (X, Δ) admits a Kähler-Einstein metric.

6 Existence theorems

A *current* on an orbifold (X, Δ) is a collection of $\text{Gal}(\varphi)$ -invariant currents on any uniformiser (U, φ) , satisfying the usual compatibility condition with respect to injections of uniformisers. In case X is smooth, orbifold differential forms on (X, Δ) are *more* than ordinary differential forms on X . By duality orbifold currents on (X, Δ) are *less* than ordinary currents on X : they are the continuous functionals on $\Lambda^k(X)$ that can be extended to the larger space $\Lambda^k(X, \Delta)$. For *positive* (p, p) -currents there is no difference between the two notions, since every positive current has measure coefficients, and every orbifold differential form has continuous coefficients. If γ is a continuous hermitian form on a compact orbifold (X, Δ) , an orbifold *Kähler current* is a closed positive (orbifold) current T of bidegree $(1, 1)$ such that for some positive constant c , $T \geq c\gamma$ in the sense of orbifold currents, that is $\langle T - c\gamma, \eta \rangle \geq 0$ for any positive $\eta \in \Lambda^{n-1, n-1}(X, \Delta)$. The definition does not depend on the choice of γ , since X is compact.

If (X, Δ) is a Fano orbifold, $G \subset \text{Aut}(X, \Delta)$ is a compact subgroup and ω is a G -invariant Kähler form in $2\pi c_1(X, \Delta)$, put

$$P_G^0(X, \Delta, \omega) = \{\chi \in C^0(X) : \omega + i\partial\bar{\partial}\chi \text{ is a Kähler orbifold current}\}.$$

Proposition 35 (a) *Any $\chi \in P_G^0(X, \Delta, \omega)$ is the C^0 -limit of a sequence $\varphi_n \in P_G(X, \Delta, \omega)$. (b) The functionals I_ω , J_ω , F_ω^0 and F_ω can be extended to $P_G^0(X, \Delta, \omega)$ and the extensions are continuous with respect to the C^0 -topology.*

(See Prop. 2.2 and 2.3 in [1].)

Lemma 36 ([1, Lemma 2.6]) *If $\pi : (X, \Delta_X) \rightarrow (Y, \Delta_Y)$ is an orbifold map between compact orbifolds, the direct image $\pi_* T$ of a Kähler current T on (X, Δ) is a Kähler current on (Y, Δ_Y) .*

Proof. First of all observe that if $f : (X, \Delta_X) \rightarrow (Y, \Delta_Y)$ is an orbifold map of degree d and $\alpha \in \Lambda^{2n}(Y, \Delta_Y)$, then $\int_X f^* \alpha = d \cdot \int_Y \alpha$. Next let γ_X and γ_Y be continuous hermitian forms on (X, Δ_X) and (Y, Δ_Y) respectively. Since $\pi^* \gamma_Y$ is continuous and γ_X is positive definite, there is $c_1 > 0$ such that $\gamma_X \geq c_1 \pi^* \gamma_Y$. If T is a Kähler current on (X, Δ) , by definition $T \geq c_2 \gamma_X$ for some $c_2 > 0$, so $T \geq c \pi^* \gamma_Y$ with $c = c_1 c_2 > 0$. We want to prove that for any positive form $\eta \in \Lambda^{n-1, n-1}(Y, \Delta_Y)$, $\langle \pi_* T, \eta \rangle \geq c \cdot \deg \pi \cdot \langle \gamma_Y, \eta \rangle$. Choose orbifold charts (V, ψ) on (Y, Δ_Y) and (U_i, φ_i) on (X, Δ) such that $\pi^{-1}(\psi(V)) = \sqcup_i \varphi_i(U_i)$. Denote by $\tilde{T}_i, \tilde{\eta}$ and $\tilde{\gamma}_Y$ the local representations in the orbifold charts and by $\tilde{\pi}_i : U_i \rightarrow V$ the liftings of π . We can assume $\text{supp}(\eta) \subset \psi(V)$. Then

$$\begin{aligned} \langle \pi_* T, \eta \rangle &= \langle T, \pi^* \eta \rangle = \sum_i \frac{\langle \tilde{T}_i, \tilde{\pi}_i^* \tilde{\eta} \rangle}{|\text{Gal}(\varphi_i)|} \geq \sum_i \frac{c \cdot \langle \tilde{\pi}_i^* \tilde{\gamma}_Y, \tilde{\pi}_i^* \tilde{\eta} \rangle}{|\text{Gal}(\varphi_i)|} = \\ &= \sum_i \frac{c}{|\text{Gal}(\varphi_i)|} \int_{U_i} \tilde{\pi}_i^* (\tilde{\gamma}_Y \wedge \tilde{\eta}) = c \cdot \left(\sum_i \frac{\deg \tilde{\pi}_i}{|\text{Gal}(\varphi_i)|} \right) \cdot \int_V (\tilde{\gamma}_Y \wedge \tilde{\eta}). \end{aligned}$$

Since

$$\sum_i \frac{\deg \tilde{\pi}_i}{|\text{Gal}(\varphi_i)|} = \frac{\deg \pi}{|\text{Gal}(\psi)|}$$

we finally get

$$\langle \pi_* T, \eta \rangle \geq c \int_{\psi(V)} (\gamma_Y \wedge \eta)$$

and this proves the lemma. $\square$

Lemma 37 ([1, Lemma 2.7]) *Let $\pi : (X, \Delta) \rightarrow (Y, \Delta_Y)$ be an orbifold map between n -dimensional Kähler orbifolds. Let ω_Y be a Kähler metric on (Y, Δ_Y) and $\chi \in P^0(Y, \Delta_Y, \omega_Y)$ a continuous potential such that $\pi^* \chi \in C^\infty(X, \Delta)$. Then*

$$F_{\pi^* \omega_Y}^0(\pi^* \chi) = F_{\omega_Y}^0(\chi). \quad (34)$$

Theorem 38 *Let (X, Δ_X) and (Y, Δ_Y) be Fano orbifolds, $\pi : (X, \Delta) \rightarrow (Y, \Delta_Y)$ an orbifold Galois covering of degree d with $G = \text{Gal}(\pi)$, ω_Y a Kähler-Einstein metric on (Y, Δ_Y) and $\omega \in 2\pi c_1(X, \Delta)$ a G -invariant Kähler metric. Assume that numerically $R^{\text{orb}}(\pi) \equiv -\beta(K_X + \Delta_X)$ for some $\beta \in \mathbb{Q}_+$. Then there is a constant C such that for any $\varphi \in P_G(X, \Delta, \omega)$*

$$F_\omega^0(\varphi) \geq \frac{1}{1+\beta} \log \left[\frac{1}{V} \int_X e^{-(1+\beta)\varphi} \pi^* \omega_Y^n \right] - C. \quad (35)$$

The proof is identical to that of Theorem 2.2 in [1] and depends on the previous lemmata. Notice that a G -invariant orbifold Kähler metric ω always exists since, according to Definition 13, $G \subset \text{Aut}(X, \Delta)$.

Theorem 39 *Let (X, Δ) , (X_1, Δ_1) , ..., (X_k, Δ_k) be n -dimensional Fano orbifolds. Assume that each (X_i, Δ_i) admits a Kähler-Einstein metric and that $\pi_i : (X, \Delta) \rightarrow (X_i, \Delta_i)$ are orbifold Galois coverings such that*

1. *the groups $\text{Gal}(\pi_i)$ are all contained in some compact subgroup of $\text{Aut}(X, \Delta)$;*
2. *$R^{\text{orb}}(\pi_i) \equiv -\beta_i(K_X + \Delta)$ for some $\beta_i \in \mathbb{Q}_+$.*

Define $\eta \in C^\infty(X, \Delta)$ by

$$\frac{1}{k} \sum_{i=1}^k \pi_i^* \omega_i^n = \eta \omega^n, \quad (36)$$

put $c := \sup\{\lambda \geq 0 : \eta^{-\lambda} \in L^1(X, \omega^n)\}$ and $\beta := \min \beta_i$. If

$$\frac{1}{c} < \beta, \quad (37)$$

then (X, Δ) admits a Kähler-Einstein metric.

The proof is the same as that of Theorem 2.3 and Proposition 2.4 in [1].

Remark 40 *If there is only one covering ($k = 1$) and X is smooth, then c is simply the complex singularity exponent (that is the log canonical threshold) of the pair (X, R^{orb}) (see [14] and [17]). On the other hand if there are enough coverings and the intersection of the ramification divisors $R^{\text{orb}}(\pi_i)$ is empty, then $c = +\infty$ and (37) is automatically satisfied.*

7 Applications

Here we exhibit some concrete examples where Theorem 39 can be used to prove the existence of Kähler-Einstein metrics on orbifolds.

Theorem 41 *Let X be a Fano manifold, $\sum_{i=1}^N D_i$ a divisor with local normal crossing and ω a Kähler-Einstein metric on X . Given integers $m_i > 1$ put $\Delta = \sum_i (1 - 1/m_i) D_i$. If $\Delta \equiv -\delta K_X$ with $\delta \in (0, 1)$ and*

$$m_i - 1 < \frac{\delta}{1 - \delta} \quad (38)$$

for any $i = 1, \dots, N$, then (X, Δ) is a Fano orbifold and has an orbifold Kähler-Einstein metric.

Proof. (X, Δ) is a Fano orbifold because $K_X + \Delta = (1 - \delta)K_X$ and $\delta < 1$. As observed in Example 15 the map $\text{id} : (X, \Delta) \rightarrow X$ is an orbifold Galois cover and we want to apply Proposition 39 to it. The ramification divisor is just $R^{\text{orb}} = \Delta$ so

$$R^{\text{orb}}(\text{id}) = -\beta(K_X + \Delta)$$

with $\beta = \delta/(1 - \delta)$. It remains to check that (38) implies (37). Let x be any point in X . Choose a system of coordinates $(V, z^1, \dots, z^n)$ on X as in Example 11 and let (U, φ) be the corresponding orbifold chart for (X, Δ) as in (6). Then on $\varphi(U) = V$

$$R^{\text{orb}} = \Delta = \sum_{j=1}^k \left(1 - \frac{1}{m'_j}\right) \{z_j = 0\} \quad (39)$$

so that in the notation of (36), $\eta(z) = \gamma(z)|f(z)|^2$ on U , where $f(z) = z_1^{m_1-1} \dots z_k^{m_k-1}$ and γ is a smooth positive function. Set $c_x = \sup\{\lambda \leq 0 : \int_U |f|^{-2\lambda} < +\infty\}$. Since

$$\int_U |f|^{-2\lambda} = \text{const} \cdot \prod_{j=1}^k \int_D |z|^{-2\lambda(m'_j-1)} \quad (40)$$

where D is the disk in $\mathbb{C}$, we get that $|f|^{-2\lambda} \in L^1_{\text{loc}}$ on U iff $\lambda < 1/(m'_j - 1)$. So $c_x = \min\{1/(m'_j - 1) : 1 \leq j \leq k\}$,

$$c = \sup_{x \in X} c_x = \min_i \frac{1}{m_i - 1} \quad (41)$$

and

$$\frac{1}{c} = \max(m_i - 1) < \frac{\delta}{1 - \delta} = \beta. \quad (42)$$

□

Example 42 Let some divisors $D_i \in |\mathcal{O}_{\mathbb{P}^n}(d_i)|$, and some integers $m_i > 1$ be given for $i = 1, \dots, N$. Let m_1 be the greatest of the m_i 's. Put $\Delta = \sum_i (1 - 1/m_i) D_i$ and

$$\delta = \frac{\sum_i d_i (1 - \frac{1}{m_i})}{n + 1}. \quad (43)$$

Assume that

1. $\sum_i D_i$ is local normal crossing;
2. $\delta < 1$;
3. $m_1(1 - \delta) < 1$.

Then the orbifold $(X, \Delta) = (X, \Delta)$ admits an orbifold Kähler-Einstein metric of positive scalar curvature.

Example 43 (Compare [10, Note 36]) Let D_i be $n + 2$ hyperplanes in general position in $\mathbb{P}^n$: $D_i = \{z_i = 0\}$ for $i = 0, \dots, n$, $D_{n+1} = \{z_0 + \dots + z_n = 0\}$. Set

$$\Delta = \sum_{i=0}^{n+1} (1 - \frac{1}{m_i}) D_i.$$

Then $(\mathbb{P}^n, \Delta)$ has an orbifold Kähler-Einstein metric as soon as

$$1 < \sum_{i=0}^{n+1} \frac{1}{m_i} < 1 + (n + 1) \min_i \frac{1}{m_i} \quad (44)$$

As in [10], many numerical examples come from Euclid's or Sylvester's sequence (cf. [23, A000058]). This is defined by the recursion relation

$$c_{k+1} = c_1 \cdots c_k + 1 = c_k^2 - c_k + 1$$

beginning with $c_1 = 2$. The sequence grows doubly exponentially, and it starts as

$$2, 3, 7, 43, 1807, 3263443, 10650056950807, \dots$$

It is easy to see that

$$\sum_{i=1}^n \frac{1}{c_i} = 1 - \frac{1}{c_{n+1} - 1} = 1 - \frac{1}{c_1 \cdots c_n}.$$

We get many new examples by taking

$$(m_0 = c_1, m_1 = c_2, \dots, m_n = c_{n+1} - 2, m_{n+1}).$$

Then

$$\sum_{i=0}^n \frac{1}{m_i} = 1 + \frac{1}{(c_{n+1} - 1)(c_{n+2} - 2)}.$$

Thus our conditions are satisfied as long as

$$c_{n+1} - 2 < m_{n+1} < n(c_{n+1} - 1)(c_{n+2} - 2)$$

and m_{n+1} is relatively prime to the other m_i .

Another case when Theorem 39 works is for degree 2 Del Pezzo surfaces S . Here we consider the case when S is allowed to have cyclic quotient singularities. These are necessarily of the form $\mathbb{C}^2/\mathbb{Z}_n$ where the group action is given by $(u, v) \mapsto (\epsilon u, \epsilon^{-1}v)$ where ϵ is a primitive n -th root of unity. The $\mathbb{Z}_n$ -invariant functions are generated by u^n, v^n, uv . This singularity is denoted by A_{n-1} .

For any degree 2 Del Pezzo surface S the anticanonical class is ample and it gives a degree 2 cover $\pi : S \rightarrow \mathbb{P}^2$. If H denotes the hyperplane class on $\mathbb{P}^2$, then $-K_S = \pi^*H$. The double cover π ramifies along a quartic curve C , thus $R = \frac{1}{2}\pi^*C = \pi^*2H$, $\beta = 2$ and to apply Theorem 39 we need to ensure that $\eta^{-\lambda}$ be integrable for $\lambda \leq \frac{1}{2}$. The singularities of π lie over the singularities of C , an A_{n-1} -singularity of S lies over an A_{n-1} -singularity of C (cf. [6, p.87]) and we can find local coordinates (x, y) on $\mathbb{P}^2$ such that S is locally isomorphic to some neighbourhood of the origin to the affine surface $\{(x, y, t) \in \mathbb{C}^3 : t^2 = x^2 + 4y^n\}$, the map π being given simply by $\pi(x, y, t) = (x, y)$. An orbifold chart is given by $\varphi : U \subset \mathbb{C}^2 \rightarrow S$ where $\varphi(u, v) = (u^n - v^n, uv, u^n + v^n)$. Thus $\varphi^*\pi^*(dx \wedge dy) = n(u^n + v^n) \cdot du \wedge dv$ and $\eta(u, v) = \text{const} \cdot |u^n + v^n|^2$. It is easy to see by direct integration or by blowing up (see e.g. [18, Prop. 6.39 p. 168]) that for $n \geq 2$, $|u^n + v^n|^{-2\lambda}$ is integrable if and only if $\lambda < \frac{2}{n}$. Thus Theorem 39 applies as long as $\frac{1}{2} < c = \frac{2}{n}$, that is for $n < 4$. This proves Theorem 2.

One can also give a different proof of the following result of Mabuchi and Mukai [19, Corollary C].

Theorem 44 *A diagonalizable singular Del Pezzo surface of degree 4 admits an orbifold Kähler-Einstein metric.*

A quartic Del Pezzo surface S is the intersection of two quadrics in $\mathbb{P}^4$, $S = Q_1 \cap Q_2$. It is said to be diagonalizable if both Q_1 and Q_2 can be put simultaneously in diagonal form. If S is singular then in suitable coordinates it is given by equations

$$h_0 := x_0^2 + x_1^2 + x_2^2 + x_3^2 + x_4^2 = 0 \quad \text{and} \quad h_1 := \lambda_2 x_2^2 + \lambda_3 x_3^2 + \lambda_4 x_4^2 = 0$$

If two of the λ_i coincide then S is a quotient of $\mathbb{P}^1 \times \mathbb{P}^1$ and so has an orbifold Kähler-Einstein metric (see [19, p.136]). Thus assume that the λ_i are distinct nonzero complex numbers. For $i = 2, 3, 4$, the equation $\lambda_i h_0 - h_1 = 0$ does not involve x_i , and by dropping the x_i variable we get smooth quadrics

$$Q_i = \{(\lambda_i h_0 - h_1 = 0)\} \subset \mathbb{P}^3.$$

The map $\pi_i : S \rightarrow Q_i$ given by forgetting x_i is a double cover ramified over the hyperplane section $S \cap \{x_i = 0\}$. Since the Q_i are smooth two-dimensional quadrics, they are Kähler-Einstein. On the other hand, the divisors $R^{orb}(\pi_i)$ are disjoint, so η is strictly positive on all S , $c = \infty$ and Theorem 39 yields that S admits an orbifold Kähler-Einstein metric.

Acknowledgments *We thank C. Arezzo, Ch. Boyer, K. Galicki and G.P. Pirola for useful comments. J.K. was partially supported by the NSF under grant number DMS-0200883.*

References

- [1] C. Arezzo, A. Ghigi, and G. P. Pirola. Symmetries, Quotients and Kähler-Einstein metrics. 2003. Preprint.
- [2] Thierry Aubin. *Some nonlinear problems in Riemannian geometry*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1998.
- [3] W. L. Baily. On the imbedding of V -manifolds in projective space. *Amer. J. Math.*, 79:403–430, 1957.
- [4] Walter L. Baily, Jr. The decomposition theorem for V -manifolds. *Amer. J. Math.*, 78:862–888, 1956.
- [5] Shigetoshi Bando and Toshiki Mabuchi. Uniqueness of Einstein Kähler metrics modulo connected group actions. In *Algebraic geometry, Sendai, 1985*, pages 11–40. North-Holland, Amsterdam, 1987.

- [6] W. Barth, C. Peters, and A. Van de Ven. *Compact complex surfaces*, volume 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*. Springer-Verlag, Berlin, 1984.
- [7] Joseph E. Borzellino. Orbifolds of maximal diameter. *Indiana Univ. Math. J.*, 42(1):37–53, 1993.
- [8] Jean Pierre Bourguignon. Métriques d’Einstein-Kähler sur les variétés de Fano: obstructions et existence (d’après Y. Matsushima, A. Futaki, S. T. Yau, A. Nadel et G. Tian), Séminaire Bourbaki, Vol. 1996/97, Astérisque, 245, 1997, 277–305
- [9] Charles P. Boyer and Krzysztof Galicki. The twistor space of a 3-Sasakian manifold. *Internat. J. Math.*, 8(1):31–60, 1997.
- [10] Charles P. Boyer, Krzysztof Galicki, and János Kollár. Einstein metrics on spheres. *arXiv:math.DG/0309408*, 2003. Preprint, to appear in *Annals of Mathematics*.
- [11] Henri Cartan. Quotient d’un espace analytique par un groupe d’automorphismes. In *Algebraic geometry and topology.*, pages 90–102. Princeton University Press, Princeton, N. J., 1957. A symposium in honor of S. Lefschetz,.
- [12] Yuan-Jen Chiang. Spectral geometry of V -manifolds and its application to harmonic maps. In *Differential geometry: partial differential equations on manifolds (Los Angeles, CA, 1990)*, volume 54 of *Proc. Sympos. Pure Math.*, pages 93–99. Amer. Math. Soc., Providence, RI, 1993.
- [13] Pierre Deligne and G. Daniel Mostow. *Commensurabilities among lattices in $PU(1, n)$* , volume 132 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1993.
- [14] Jean-Pierre Demailly and János Kollár. Semi-continuity of complex singularity exponents and Kähler-Einstein metrics on Fano orbifolds. *Ann. Sci. École Norm. Sup. (4)*, 34(4):525–556, 2001.
- [15] Hans Grauert and Reinhold Remmert. *Coherent analytic sheaves*, volume 265 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1984.
- [16] Shoshichi Kobayashi, *Topology of positively pinched Kaehler manifolds*, Tôhoku Math. J. (2) **15** (1963), 121–139.

- [17] János Kollár. Singularities of pairs. In *Algebraic geometry—Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, pages 221–287. Amer. Math. Soc., Providence, RI, 1997.
- [18] János Kollár and Karen E. Smith, and Alessio Corti. *Rational and nearly rational varieties*, Cambridge University Press, Cambridge, 2004.
- [19] Toshiaki Mabuchi and Shigeru Mukai. Stability and Einstein-Kähler metric of a quartic del Pezzo surface, in *Einstein metrics and Yang-Mills connections (Sanda, 1990)*, volume 145 of *Lecture Notes in Pure and Appl. Math.*, pages 133–160, Dekker, New York, 1993
- [20] Yasuhiro Nakagawa. An isoperimetric inequality for orbifolds. *Osaka J. Math.*, 30(4):733–739, 1993.
- [21] Peter Orlik and Philip Wagreich, Seifert n -manifolds, *Invent. Math.*, 28 (1975), 137–159.
- [22] David Prill. Local classification of quotients of complex manifolds by discontinuous groups. *Duke Math. J.*, 34:375–386, 1967.
- [23] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, AT&T Research, <http://www.research.att.com/~njas/sequences/>, 2003.
- [24] Gang Tian. On Kähler-Einstein metrics on certain Kähler manifolds with $C_1(M) > 0$. *Invent. Math.*, 89(2):225–246, 1987.
- [25] Gang Tian. On Calabi’s conjecture for complex surfaces with positive first Chern class, *Invent. Math.*, 101, 1990, 101–172.
- [26] Gang Tian. Kähler-Einstein metrics with positive scalar curvature. *Invent. Math.*, 130(1):1–37, 1997.
- [27] Gang Tian. *Canonical metrics in Kähler geometry*. Birkhäuser Verlag, Basel, 2000. Notes taken by Meike Akveld.
- [28] Xu-Jia Wang and Xiaohua Zhu. Kähler-Ricci solitons on toric manifolds with positive first Chern class. *Adv. Math.*, 188(1):87–103, 2004.

Università di Milano Bicocca

`alessandro.ghigi@unimib.it`

Princeton University, Princeton NJ 08544-1000

`kollar@math.princeton.edu`